

# Mixed Layer data set of Argo, Grid Point Value: MILA GPV ver. 1.0

2017.03.30

Japan Agency for Marine-Earth Science and Technology (JAMSTEC)

## Introduction

JAMSTEC produces a global data set of gridded mixed layer depth (MLD) using Argo temperature and salinity profiles, named "MILA GPV (Mixed Layer data set of Argo, Grid Point Value)". Since quality of the profile data which covered fully in the ice-free global ocean is high and uniform and vertical resolution of the profiles is fine (at least 5 dbar in the surface layer), error of MLD is expected to be smaller. Although the coverage of Argo float array was not good in the earlier stage of Argo project, observing floats almost covered in the World Ocean after 2004. The Argo quality controlled profile data (real time QC and/or delayed mode QC) is obtained from Argo Global Data Assembly Center (GDAC). For the obtained data, further quality control is carried out to avoid including inappropriate profile data. To make MLD data set, the finite difference method is used based on sea surface temperature and sea surface potential density. Among the values in the difference of temperature and potential density, mixed layer is chosen with the minimum value between the two values. Two pairs of the temperature and potential density criteria are applied, which are generally used in a lot of researches. The grid point value of MLD is calculated using a median after estimated MLD for individual profiles. Users can use the two data set for their aim of research.

## Method and dataset information

Before estimating MLD, quality controls are carried out for Argo profile data which are quality controlled real time and/or delayed mode data. The quality controls are 1) to check temperature and density inversion, 2) to confirm existence of surface values above 15 dbar and 3) to remove over 3 times of standard deviation at each levels using World Ocean Atlas 2005 (Antonov et al., 2006; Locarnini et al., 2006). For individual profiles, vertical interpolation based on Akima method is done for each 1 dbar (Akima, 1970). Using the interpolated profiles, MLD, isothermal layer thickness (TLD), isopycnal thickness (PLD) and averaged temperature and salinity in the MLD are estimated for each profile. The MLD is defined with the minimum value between TLD and PLD. The TLD and PLD are calculated using the finite difference method based on the temperature and potential density values at 10 dbar. The two pairs of criteria are chosen, which are used in recent research, that is, 1) 0.5°C of difference in temperature ( $\Delta T$ ) and 0.125 kg m<sup>-3</sup> of difference in potential density ( $\Delta \sigma_\theta$ ; e.g.,

Monteley and Levitus,1997) and 2) 0.2 °C of  $\Delta T$  and 0.03 kg m<sup>-3</sup> of  $\Delta\sigma_\theta$  (e.g., de Boyer Montégut et al., 2004). MLD and the other parameters are gridded in 2°x2° boxes, whose values are calculated using median at each grid; however averaging is done in the case if the number of data within grid is less than 4. The data set is made in the ice-free global ocean a) for each 10 days from 2001- on going, b) monthly average from 2001 to 2010, c) for one month from 2001 – on going. The data set will be renewed within one or two months. The detailed of method, quality control and data set are described in Hosoda et al. (2010).

(1) Parameters and criteria for each gridded data set in a)- c)

- i) Mixed layer depth
  - i-1) 0.03kg m<sup>-3</sup> for  $\Delta\sigma_\theta$  and 0.2°C for  $\Delta T$
  - i-2) 0.125kg m<sup>-3</sup> for  $\Delta\sigma_\theta$  and 0.5°C for  $\Delta T$
- ii) Isothermal layer thickness
  - ii-1) 0.2°C for  $\Delta T$
  - ii-2) 0.5°C for  $\Delta T$
- iii) Isopycnal layer thickness
  - iii-1) 0.03kg m<sup>-3</sup> for  $\Delta\sigma_\theta$
  - iii-2) 0.03kg m<sup>-3</sup> for  $\Delta\sigma_\theta$
- iv) Averaged temperature in MLD
- v) Averaged salinity in MLD
- vi) Averaged potential density in MLD
- vii) The number of profiles in each grid

(2) File name

- i) Monthly mean mixed layer
  - File name: mila\_[mld/mld2/05dC/02dC/0125st/003st]<sup>(a)</sup>[mld/tmp/sal/pod]  
<sup>(b)</sup>\_[yyyyymm]<sup>(c)</sup>00\_010.nc
  - (a) “mld”: mixed layer depth ((1) i-1), “mld2”: mixed layer depth ((1) i-2), “02dC”: isothermal layer depth ((1) ii-1), “05dC”: isothermal layer depth: ((1) ii-2), “003st”: isopycnal layer depth ((1) iii-1), “0125st”: isopycnal layer depth ((1) iii-2)
  - (b) “mld”: mixed layer depth, “tmp”: averaged temperature, “sal”: averaged salinity, “pod”: averaged potential density
  - (c) “yyyy”: year, “mm”: month
  - Period: January 1, 2001~

ii) 10-day mean mixed layer

File name: mila10d\_[mld/mld2/05dC/02dC/0125st/003st]<sup>(a)</sup>[mld/tmp/sal/pod]

<sup>(b)</sup>\_[yyyyymmdd] <sup>(c)</sup>\_010.nc

(a) “mld”: mixed layer depth ((1) i-1), “mld2”: mixed layer depth ((1) i-2), “02dC”: isothermal layer depth ((1) ii-1), “05dC”: isothermal layer depth: ((1) ii-2), “003st”: isopycnal layer depth ((1) iii-1), “0125st”: isopycnal layer depth ((1) iii-2)

(b) “mld”: mixed layer depth, “tmp”: averaged temperature, “sal”: averaged salinity, “pod”: averaged potential density

(c) “yyyy”: year, “mm”: month, “dd”: first day of 10-day average

Period: January 1-10, 2001~

iii) Monthly mean mixed layer

File name: milaclim\_[mld/mld2/05dC/02dC/0125st/003st]<sup>(a)</sup>[mld/tmp/sal/pod]

<sup>(b)</sup>\_[mm] <sup>(c)</sup>\_010.nc

(a) “mld”: mixed layer depth ((1) i-1), “mld2”: mixed layer depth ((1) i-2), “02dC”: isothermal layer depth ((1) ii-1), “05dC”: isothermal layer depth: ((1) ii-2), “003st”: isopycnal layer depth ((1) iii-1), “0125st”: isopycnal layer depth ((1) iii-2)

(b) “mld”: mixed layer depth, “tmp”: averaged temperature, “sal”: averaged salinity, “pod”: averaged potential density

(c) “mm”: month

Period: January~ December

(3) Area and spatial resolution of dataset, and definition of “TIME” variable

Spatial area of MILA GPV is [74.5°S-79.5°N, 0.5°-359.5°]. Horizontal grid size is 1° x1°, which is interpolated from original analysis data (2° x2°). “TIME” variable in each file is defined as month from Jan.1, 1990 for (2) i), day from Jan.1, 1990 for (2) ii) and month for (2) iii).

(4) Data format

Format of MILA GPV is described with Network Common Data Form (NetCDF) on the basis of COARDS rule. The NetCDF data can convert to ascii format by using “ncdump” software. If some analysis software such as “GrADS” and “Ocean Data View(ODV)” are used, NetCDF is also available.

## About this document

Since this document will be occasionally renewed, the date and version of the document must be checked.

## Usage of MILA GPV v1.0

- Reprint without permission, the re-distribution, the modification, and the commercial use are prohibited though JAMSTEC does not disturb the free use of this data in principle.

Hosoda, S., T. Ohira, K. Sato, and T. Suga (2010): Improved description of global mixed-layer depth using Argo profiling floats, <i>J. Oceanogr</i> , 66, 2010,773-787.
---

- Please source the above when you make the result using this data public.
- JAMSTEC doesn't assume the responsibility to any damage of the user of this data.
- Documents about MOAA GPV in the web site are occasionally modified or corrected. Users should check the version when any research result from the dataset is published.

## Update History of this document for MILA GPV v1.0

First Edition: 30<sup>th</sup> March, 2017

## References

- Akima, H. (1970): A new method for interpolation and smooth curve fitting based on local procedures. *J. Assoc. Comput. Mech.*, **17**, 589– 603.
- Antonov, J. A., R. A. Locarnini, T. P. Boyer, H. E. Garcia, and A. Mishonov (2006): World Ocean Atlas 2005, Vol. 2: Salinity. Ed. S. Levitus. NOAA Atlas NESDIS 62, U.S. Government Printing Office, Wash, D.C.
- Argo (2000): Argo float data and metadata from Global Data Assembly Centre (Argo GDAC), Ifremer, <http://dx.doi.org/10.12770/1282383d-9b35-4eaa-a9d6-4b0c24c0cfc9>.

- de Boyer Montégut, C., G. Madec, A. S. Fischer, A. Lazar, and D. Iudicone (2004): Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology. *J. Geophys. Res.*, **109**, C12003, doi:10.1029/2004JC002378.
- Hosoda, S., T. Ohira, K. Sato, and T. Suga (2010): Improved description of global mixed-layer depth using Argo profiling floats, *J. Oceanogr*, 66, 2010,773-787.
- Locarnini, R. A., A. Mishonov, Antonov, J. A., T. P. Boyer, and H. E. Garcia (2006): World Ocean Atlas 2005, Vol. 1: Temperature. Ed. S. Levitus. NOAA Atlas NESDIS 61, U.S. Government Printing Office, Wash, D.C.
- Monterey, G. and S. Levitus (1997): *Seasonal Variability of Mixed Layer Depth for the World Ocean*. NOAA Atlas NESDIS 14, 100 pp.