

Cloning and overexpression of the *rpoZ* gene encoding RNA polymerase ω subunit from a deep-sea piezophilic bacterium, *Shewanella violacea* strain DSS12

Hiroaki KAWANO*^{1,2} Yasuo SUZAKI*¹ Junko FUKUCHI*² Kaoru NAKASONE*³ Fumiyoshi ABE*²
Chiaki KATO*⁴ Yasuhiko YOSHIDA*¹ Ron USAMI*¹ Koki HORIKOSHI*^{1,2}

We have cloned the *rpoZ* gene, encoding RNA polymerase ω protein, by PCR approach from the deep-sea piezophilic and psychrophilic bacterium, *Shewanella violacea* strain DSS12. The cloned gene, 285 bp in length, was found to encode a protein consisting of 93 amino acid residues with a molecular mass of 10,227 Da. Significant homology was evident comparing the *rpoZ* protein of *S. violacea* with that of *Escherichia coli* K-12 (71% identity), *Vibrio cholerae* (70% identity) and *Haemophilus influenzae* (61% identity). Phylogenetic analysis of RpoZ proteins of several bacteria suggested that *S. violacea* is independent from other bacteria. We constructed expression plasmid to overproduce the RpoZ protein and transformed into *E. coli* JM109 as a host of overproduction. Upon induction, the recombinant protein encoded by plasmid pQrpoZ was overexpressed and purified using Ni²⁺ affinity column. The result of SDS-PAGE during purification suggests that a chimeric RNA polymerase ($\alpha_2\beta\beta'\sigma^{\omega}$) was assembled in *E. coli*.

Keywords : *Shewanella violacea*, RNA polymerase ω protein, *rpoZ* gene, expression plasmid

*1 Department of Applied Chemistry, Faculty of Engineering, Toyo University

*2 The DEEP STAR Group, Japan Marine Science and Technology Center

*3 Department of Chemistry and Environmental Technology, School of Engineering, Kinki University

*4 Marine Ecosystems Research Department, Japan Marine Science and Technology Center

1. Introduction

The moderately piezophilic and psychrophilic bacterium *Shewanella violacea* strain DSS12 from Ryukyu Trench (depth 5,110 m) grows optimally at 30 MPa and 8°C, but also grows at atmospheric pressure (0.1 MPa) and 8°C [1, 2]. These growth properties are useful for comparative study of cell physiology under high and low pressure conditions.

Recently an operon identified as a pressure-regulated operon, with a promoter activated under high-pressure conditions, was cloned and characterized from this strain [3-5]. We have reported that gene expression from this operon is controlled at the transcriptional level by elevated pressure [3]. However, the mechanisms controlling transcription of this operon at the molecular level remain to be elucidated. One approach to understanding the basis of gene expression under defined condition is via detailed characterization of transcriptional machinery involved.

Thus, cloning of the genes for RNA polymerase subunits from piezophiles and the establishment of an *in vitro* system for reconstitution of the enzyme are essential. Transcription in eubacteria is mediated by an RNA polymerase, which governs the selectivity of the promoter sequences of a gene. It is a multisubunit complex composed of mainly α , β , β' and one of several σ subunits [6]. Among these subunits, we have previously reported piezoresponse of *S. violacea* ω operon and some positive regulators may exist and interact with RNA polymerase [7]. It is essential to elucidate and to characterize several factors interact with RNA polymerase and relate to transcriptional regulation at several conditions. In the case of *E. coli*, one of these factors named ω protein is present in both highly purified core and holo RNA polymerase, but the roles of this protein is very little known. One of estimated functions of *E. coli* ω protein is a regulator of RNA polymerase, to control promoter selectivity at specific stringent promoters, *rpsA* and *rplJ*, mediated by ppGpp [8]. In this study, we report the isolation of the *rpoZ* gene encoding RNA polymerase ω protein from *S. violacea*. The results of phylogenetic analyses based on comparison of the RpoZ protein, construction of expression plasmid, and its affinity purification will be described.

2. Materials and Methods

2.1. Cloning of the *rpoZ* gene from *S. violacea*

Two degenerate oligonucleotides were designed and synthesized, based on highly conserved sequences, MARVTV and VIALREI, in RpoZ proteins. The nucleotide sequences of oligonucleotides were 5'-ATGGCNCNG-TAACWGTN-3' (primer 1) and 5'-DATYTCRCGYAG-NGCDATWAC-3' (primer 2). The PCR product obtained using *S. violacea* chromosomal DNA as the template and the above primers, was cloned into the pCR2.1 vector (Invitrogen Co., Carlsbad, CA) and the nucleotide sequence was determined by dye terminator method using a DNA sequencer model 377 (Perkin-Elmer/applied Biosystems Co., Foster, CA). Based on the determined nucleotide sequence, two other PCR primers were designed for use in amplification of the digoxigenin (DIG)-labeled *rpoZ* fragment. The DIG-labeled fragment was employed as the probe for plaque hybridization in screening by means of the DIG detection system (Boehringer Mannheim Co., Mannheim, Germany).

2.2. Southern blotting analysis and mapping of the *rpoZ* gene of *S. violacea*

The *rpoZ* gene was identified from λ phage library of *S. violacea* chromosomal DNA and significant homologies to known *rpoZ* gene. Based on the determined nucleotide open reading frame sequence, two other PCR primers were designed for use in amplification of the digoxigenin (DIG)-labeled *rpoZ* fragment and for use in cloning (method of 2.5.). The nucleotide sequences of the oligonucleotides were 5'-ATGGCTCGCGTAACTGTAGAA-3'(primer 1) and 5'-TTTATAATATGTCGTTGC-GACC-3' (primer 2). The DIG labeled fragment was employed as the probe for southern blotting analysis. The chromosomal DNA of *S. violacea* was digested with *Bam*H I, *Eco*R I, *Hind* III, *Sal* I and *Pst* I, digested fragments were separated on 1% agarose gel and blotted onto a PVDF membrane. The fragments were detected by means of the DIG detection system (Boehringer Mannheim Co., Mannheim, Germany).

2.3. Construction of a λ phage library of the chromosome of *S. violacea* and screening of the library for the *rpoZ* gene region

Chromosomal DNA isolated from *S. violacea* was partially digested with *Sau*3AI. These fragments were

inserted into the *Bam*H I site of lambda DASH II (Stratagene Co., La Jolla, CA). Then, *in vitro* packaging of the ligated DNA was performed using GIGAPACK III XL packaging extracts (Stratagene Co.) according to the manufacture's instructions. The λ phage library was screened for plaque hybridization with the *rpoZ* probe and several positive clones were obtained. The positive clones containing the *rpoZ* gene were each purified by several single plaque isolation steps. Each of the inserts in λ phage was amplified by long PCR and was sub-cloned into the pCR-Blunt vector (Invitrogen Co.). For sequencing of these cloned fragments, the random shotgun sequencing method was used with a DNA sequencer model 377 (Perkin-Elmer/applied Biosystems Co.). Assembling and editing of the determined DNA sequences were performed with AutoAssembler Version 2.0 (Perkin-Elmer/Applied Biosystems Co.). GENE-TYX-MAC version 10.1 from software Development (Tokyo, Japan) was used for sequence analysis.

2.4. Phylogenetic analysis of the RpoZ protein

To determine the phylogenetic position of the *S. violacea* RpoZ protein, evolutionary distance was calculated and a phylogenetic tree was constructed. The alignment and calculation were performed using CLASTALX and the PHYLIP (ver. 3.573; obtained from J. Gelsenstein, University of Washington, seattle).

2.5. Construction of expression plasmid harboring *rpoZ* gene

To construct plasmids for expression of hexahistidine-tagged derivative of the RpoZ protein from strain DSS12, PCR was performed to amplify the *rpoZ* gene. The PCR product was cloned into the pCR2.1 vector and sequenced. The resulting fragment, digested with both *Bam*H I and *Hind* III, was cloned into expression plasmid pQE80L (QIAGEN) and then it was designated as pQrpoZ (Fig. 4A).

2.6. Overexpression and affinity purification of the ω subunit (RpoZ protein)

The expression plasmids of pQrpoZ were transformed into *E. coli* JM109. Overexpression of the protein was carried out by the procedure described by Nakasone et al. [7]. Overexpressed protein was fractionated by 15% SDS-PAGE and the molecular mass was determined (Fig. 4B).

3. Results and discussion


3.1. Isolation of the *rpoZ* gene of *S. violacea*

To clone a part of the *rpoZ* gene from *S. violacea*, the amino acid sequences of the RpoZ proteins from several bacteria were aligned and a highly conserved region was found (data not shown). These conserved sequences were used to design and synthesize degenerate oligonucleotides in order to amplify a part of the *rpoZ* gene from *S. violacea*. A fragment containing part of the *rpoZ* gene, approximately 170 bp, amplified by PCR with the primers, was cloned into the pCR2.1 vector and its nucleotide sequence was determined. The deduced amino acid sequence of the cloned gene was significantly similar to the RpoZ proteins of other bacteria (data not shown). To clone the complete *rpoZ* gene, the partial *rpoZ* gene fragment was labeled with DIG (digoxigenin) by PCR as a hybridization probe. From the Southern blotting analysis using DIG-labeled *S. violacea rpoZ* gene as a probe, a positive band was detected from digested fragments of *S. violacea* Chromosomal DNA with several restriction enzymes (Fig. 1A). As shown in Fig. 1A it showed that the *rpoZ* gene was present in a 36.5 kb (*Bam*H I), 8.2 kb (*Hind* III), 9.9 kb (*Eco*R I), 2.0 kb (*Pst* I) and 20.1 kb fragment (*Sal* I).

3.2. Structural analysis of the *rpoZ* gene of *S. violacea*

A λ phage library of *S. violacea* chromosomal DNA was constructed and screened using the *rpoZ* probe. Several positive clones were isolated from among 104 plaques and purified by repeated single plaque isolation. These clones were confirmed to contain the *rpoZ* gene by PCR at each isolation step. As a result of screening, we obtained several positive clones and the nucleotide sequence of one of these clones containing a DNA insert approximately 10 kb in size was determined by the random shotgun sequencing method [8]. Based on the results of sequence and Southern blotting analysis, we constructed the restriction map of *rpoZ* gene. As shown in Fig. 1B, the 10 kb DNA fragment, including *rpoZ* gene, had ten open reading frames (ORFs) and each ORFs protein names were determined with Genbank database. Besides of the *rpoZ* gene, the determined 10kb DNA fragment contained nine other genes, *pgtB*, *mdsC*, *functionally unknown gene 1*, *gmkc*, *spoT*, *functionally unknown gene 2*, *spoU*, *ribD* and *functionally unknown gene 3*, respectively. It was reported that the *rpoZ* gene of *E. coli* is located in an operon that also encodes the

(A)



(B)




Fig. 1 Results of southern blotting analysis, restriction map of the RNA polymerase ω subunit gene of *S. violacea* strain DSS12.

(A) Results of southern blotting analysis of the *rpoZ* gene in *S. violacea*.

(B) Restriction map of the *S. violacea rpoZ* 10 kb fragment containing the genes *pgtB*, *mdsC*, *unknown*, *gmkc*, *rpoZ*, *spoT*, *unknown*, *spoU*, *ribD*, *unknown*.

spoT gene whose product is involved in the metabolism of ppGpp [9-10]. The *rpoZ* gene of *S. violacea* was located at the upstream region of *spoT* gene as well as *E. coli*. From the alignment of the upstream region of *S. violacea* with that of *E. coli*, two promoter-like sequences of RNA polymerase were identified (data not shown). From these results, the transcriptional regulation of this region may be conserved in two species and raised possibility of the ω is required for transcriptional regulation by ppGpp. The open reading frame of *rpoZ* gene, 285 bp in length, was found to encode a polypeptide consisting of 94 amino acid residues with a molecular mass of 10,227 Da (Fig. 2A). Analysis of the deduced amino acid sequence confirmed that the cloned gene contained the complete sequence of the RpoZ protein (Fig. 1B). From the comparison of RpoZ protein with *Escherichia coli*

and *Vibrio cholerae*, the amino acid sequences of *S. violacea* RpoZ shows a high degree of similarity to those of RpoZ homologues (Fig. 2B). The N-terminal two-thirds of RpoZ protein is significantly conserved, but the C-terminal region is not so conserved among these organisms. The result suggests that the two-thirds of N-terminal region of RpoZ protein is important to bind with RNA polymerase molecules. As shown in Fig. 2B, it was reported the N-terminal 24 and 25 residues (Arg-Arg) of *E. coli* ω protein is efficiently cleaved by the OmpT protease [11]. This site is observed in the case of ω protein of *S. violacea* and *V. cholerae*. This feature indicating the ω protein of *S. violacea* is cleaved if it was overproduced in *E. coli* JM109 as a host. As predicted, one other fragment was observed after overproduced RpoZ protein by SDS-PAGE (described in chapter 3. 3).




Fig. 2 Nucleotide and deduced amino acid sequences of the RNA polymerase ω subunit gene of *S. violacea* strain DSS12. Nucleotide and sequence encoded by the gene is shown in single-letter notation below the codon; an in-frame stop codon is indicated by an asterisk. The numbers of the right refer to nucleotide positions. The nucleotide sequences corresponding to the PCR primers used (Primer 1 and Primer 2) are indicated by arrows. The predicted OmpT protease dependent residues are squared.

3.2. Phylogenetic analysis of the evolutionary distance of *S. violacea* RpoZ protein

From the alignment of amino acid residues of the *S. violacea* RpoZ protein with other organisms, the RpoZ protein is conserved in a wide range of organisms (data not shown). To confirm the phylogenetic position of the *S. violacea* RpoZ protein, the evolutionary distance was calculated based on comparison of 17 organisms and a phylogenetic tree was constructed (Fig. 3). The RpoZ amino acid sequences were aligned and the tree was constructed by the neighbor-joining method. The results obtained from calculation of the evolutionary distance comparing the RpoZ amino acid sequences of 17 organisms showed that the RpoZ protein of *S. violacea* is independent of other 17 organisms.

3.3. Construction of expression plasmid and overproduction of *S. violacea* ω protein

A recombinant expression plasmid harboring the *S. violacea* *rpoZ* gene was constructed (pQrpoZ) and

transformed into *E. coli* JM109 (Fig. 4B). Upon induction, the recombinant RpoZ protein was overexpressed as shown in lane 3 of Fig. 4B. As discussed in chapter 3. 2, the amino acid sequence with *E. coli* and *S. violacea*, the Arg-Arg recognition site of OmpT protease was identified. In this experiment, we detected a fragment, observed at smaller size of hexahistidine-tagged protein after induction. This fragment was probably two-thirds of ω protein. These results indicating that the *E. coli* OmpT protease can cut *S. violacea* ω protein off as well as that of *E. coli*. Hexahistidine-tagged ω protein was purified by a simple chromatographic step and the molecular mass was determined to be 14.8 kDa by SDS-PAGE (lane 4 of Fig. 4B). We purified this ω protein from soluble fraction not only for purification by nondenaturing treatments but also for detection of several interacted protein with ω protein in *E. coli*. As shown in lane 6 of Fig. 4B, we identified several protein bands fractionated simultaneously during elution steps by SDS-PAGE. In these protein bands, we identified




Fig. 3 Phylogenetic tree inferred by neighbor joining analysis of RpoZ protein sequences.




Fig. 4 Construction of expression plasmid harboring *rpoZ* gene and purification of hexahistidine-tagged ω protein.
 (A) Construction of expression plasmid
 (B) SDS-PAGE analysis of the *S. violacea* ω hexahistidine-tagged fusion protein expressed in *E. coli* JM109. Lane 1, the sizes of the molecular mass standards indicated in kDa; lane 2, before induction of expression of the *rpoZ* gene encoded by pQrpoZ; lane 3, soluble fraction, after overexpression of the *rpoZ* gene encoded by pQrpoZ was induced by treatment with 1 mM IPTG; lane 4, purified His6- ω protein from soluble fraction; lane 5, *E. coli* RNA polymerase; lane 6, fraction No.1 during elution of ω protein. An arrow indicates the position of the protein, His6- ω with a mass of 14.8 kDa, a fragment, and RNA polymerase subunits (α , β , β' , σ^{70}) of *E. coli*.

some proteins that are estimated as RNA polymerase subunits (α , β , β' , σ^{70}). This result suggests that, a chimeric RNA polymerase composed of α_2 , β , β' , σ^{70} from *E. coli* and ω from *S. violacea* was assembled in the *E. coli* JM109 harboring pQrpoZ. In addition, we also used *E. coli* TOP10 as a host of overproduction, but could not observe expression of pQrpoZ (data not shown). Using highly purified RpoZ protein, to investigate its effect on in vitro transcription process is expected. From the establishment of this overproduction method, we could obtain several recombinant ω proteins and could introduce mutation to study the role of this protein using knockout mutant.

References

- [1] Kato C, Sato T, Horikoshi K. Isolation and properties of barophilic and barotolerant bacteria from deep-sea mud samples: Biodiv. Conserv 4: 1-9. (1995)
- [2] Nogi Y, Kato C, Horikoshi K. Taxonomic studies of deep-sea barophilic *Shewanella* species, and *Shewanella violacea* sp. nov., a new barophilic bacterial species: Arch Microbiol 170: 331-338. (1998)
- [3] Kato C, Ikegami A, Usami R, Horikoshi K. Structure of genes in a pressure-regulated operon and adjacent regions from a barotolerant bacterium strain DSS12: J Mar Biotechnol 5: 210-218. (1997)
- [4] Nakasone K, Ikegami A, Kato C, Usami R, Horikoshi K. Analysis of cis-elements upstream of the pressure-regulated operon in the barophilic bacterium *Shewanella violacea* strain DSS12: FEMS Microbiol Lett 1491: 315-320. (2000)
- [5] Ikegami A, Nakasone K, Fujita M, Fujii S, Kato C, Usami R, Horikoshi K. Cloning and characterization of the gene encoding RNA polymerase sigma factor σ^{54} of deep-sea piezophilic *Shewanella violacea*: Biochim Biophys Acta 1491: 315-320. (2000)
- [6] Yura T, Ishihama A. Genetics of bacterial RNA polymerases: Annu Rev Genet. 13: 59-97. (1979)
- [7] Nakasone K, Ikegami A, Fujita M, Fujii S, Kato C, Horikoshi K. Isolation and piezoresponse of the *rpoA* gene encoding RNA polymerase *a* subunit from the deep-sea piezophilic *Shewanella violacea*: FEMS microbiol Lett 9717: 1-8. (2000)
- [8] Yamada M, Nakasone K, Tamegai H, Kato C, Usami R, Horikoshi K. Pressure regulation of soluble cytochromes *c* in a deep-sea piezophilic bacterium, *Shewanella violacea*: J. Bacteriol 182: 2945-2952. (2000)
- [9] Daniel R Gentry, Richard R Burgess. *rpoZ*, encoding the omega subunit of *Escherichia coli* RNA polymerase, is in the same operon as *spoT*: J Bacteriol 171: 1271-1277. (1989)
- [10] Igarashi K, Fujita N, Ishihama A. Promoter selectivity of *Escherichia coli* RNA polymerase: Omega factor is responsible for the ppGpp sensitivity Protein: Nucleic Acids Res 17: 81-86. (1989)
- [11] Daniel R Gentry, Richard R Burgess. Overproduction and Purification of the ω Subunit of *Escherichia coli* RNA polymerase: Protein Expr Purif 1: 81-86. (1990)

(Manuscript received 13 December 2002)