

Studies of the physiology of chemosynthetic fauna at Cold Seeps in Sagami Bay, Japan

J. P. BARRY*¹ R. E. KOCHVAR*² J. HASHIMOTO*³
Y. FUJIWARA*³ K. FUJIKURA*³ H. G. GREENE*⁴

Comparisons of cold seep habitats in Sagami Bay, Japan, and Monterey Bay, California, U. S. A. have included studies of the geology and geochemistry of seeps as well as the biology and ecology of seep-associated fauna. In this paper we present results of research concerning the physiology of chemosynthetic fauna found at cold seeps in Sagami Bay.

Haemolymph sulfide binding properties were measured for individuals of *Bathymodiolus platifrons*, *Calyptogenia soyozaki*, and an undescribed vestimentiferan worm (*Lamellibrachia* sp.) that inhabit seeps in Sagami Bay. *C. soyozaki* and *Lamellibrachia* sp. haemolymph was shown to bind sulfide more than 10 times ambient levels. Haemolymph from *B. platifrons* did not show evidence of sulfide binding at any level of ambient sulfide concentrations.

Key words : Cold seep community, Physiology, Ecology, Sagami Bay, Monterey Bay

* 1 Monterey Bay Aquarium Research Institute

* 2 Monterey Bay Aquarium

* 3 Deep Sea Research Department, Japan Marine Science and Technology Center

* 4 Moss Landing Marine Laboratories

1. Introduction

Chemosynthetic communities associated with seepage of sulfide or methane-rich fluids have been found in various settings since their discovery in the late 1970s and early 1980s (Lonsdale, 1979; Paull *et al.*, 1984; Kennicutt *et al.*, 1985) and appear to be common along continental margins (Barry *et al.*, in press a). Chemosynthetic communities dominated by vesicomyid clams, mytilid mussels, vestimentiferan or pogonophoran worms, or free-living bacteria, have been reported from hydrocarbon seeps (Kennicutt *et al.*, 1985, 1988, 1989), groundwater seeps (Paull *et al.*, 1984; Hecker, 1985), methane seeps (Kennicutt *et al.*, 1985; Dando *et al.*, 1991), accretionary prisms associated with subduction zones, other sites of tectonically-compressed sediment (Suess *et al.*, 1985; Laubier *et al.*, 1986; Boulègue *et al.*, 1987; Cadet *et al.*, 1987; Le Pichon *et al.*, 1987; Ohta and Laubier, 1987; Hashimoto *et al.*, 1987, 1989; Jollivet *et al.*, 1990; Barry *et al.*, in press a), and relict organic material buries in debris flow deposits (Embley *et al.*, 1990). Fossil cold seeps and associated faunal assemblages have also been reported from several sites (Beauchamp *et al.*, 1989; von Bitter *et al.*, 1990, 1992; Campbell and Bottjer, 1993; Aharon, 1994; Clari *et al.*, 1994).

During the past few years, comparisons of cold seep communities in Sagami Bay, Japan, and Monterey Bay, California, U. S. A. have been performed by scientists from JAMSTEC, Monterey Bay Aquarium Research Institute (MBARI), and Moss Landing Marine Laboratories (MLML). Dives at cold seeps on either side of the Pacific have been made using manned submersibles ("Shinkai 2000") and ROVs ("Dolphin 3K", Ventana), to support studies of biology, geology and geochemistry.

Similarities between the geology and geochemistry of Sagami and Monterey Bay seeps (Greene *et al.*, ms this issue), appear responsible for partial overlap in the composition of chemosynthetic fauna. Vesicomyid clams are the dominant metazoan fauna of cold seeps in both Sagami Bay (Hashimoto *et al.*, 1989) and Monterey Bay (Barry *et al.*, in press a). However, studies of the fluid chemistry in both areas show that although there is overlap among sulfide concentrations within aggregations of vesicomyid clams on either side of the Pacific (Hashimo-

to *et al.*, 1995; Barry *et al.*, in press b), the levels of sulfide and methane vary considerably among sites.

Relatively little information has been published concerning the biology of vesicomyids from cold seep habitats. Most information concerning vesicomyid biology has come from studies of *Calyptogena magnifica*, the very large species common at hydrothermal vents (Boss and Turner, 1980; see Childress and Fisher, 1992).

Research described in this paper concerns the sulfide-binding affinity of blood serum from *Calyptogena soyoe*, mytilid mussels (*Bathymodiolus platifrons*), and an undescribed vestimentiferan worm (*Lamellibrachia* sp.). Results from these studies in Sagami Bay and similar studies of cold seep vesicomyids from Monterey Bay (and hydrothermal vent sites), represent the first such comparisons of the physiology of vesicomyid clams among seep and vent sites.

2. Method

2.1 Study Site

Hashimoto *et al.* (1989) present a thorough description of cold seeps in Sagami Bay and the biota inhabiting these seeps. Geochemistry of pore waters was examined by Masuzawa *et al.* (1995), and the relation of pore fluid chemistry to the distribution of *Calyptogena soyoe* populations at the Hatsushima site cold seeps was discussed by Hashimoto *et al.* (1995). Additional information on the existence of thermal anomalies related to volcanogenic heating is presented in Naka *et al.* (1991).

2.2 Haemolymph Sulfide Binding

Sulfide binding studies were carried out on haemolymph of vesicomyid clams (*Calyptogena soyoe*; collected on "Shinkai 2000" dives 828 and 829), mytilid mussels (*Bathymodiolus platifrons*; collected on dives 829 and 831), and vestimentiferan tube worms (*Lamellibrachia* sp.; collected on dive 832). Haemolymph samples were removed from specimens immediately following recovery of the submersible. A 0.5 ml aliquot of haemolymph was immediately placed in a vial containing 0.5 ml fixing solution (1 M zinc acetate) following dissection and exsanguination of each specimen. Additional 0.5 ml aliquots were injected into dialysis cassettes (Pierce, 10k MWCO Slide-A-Lyzer), which were distributed among dialysate solutions. Sulfide stock solu-

tion was prepared by dissolving sodium sulfide crystals in nitrogen-bubbled (oxygen-free) sea water, and adjusting the pH with HCl to 8. Sulfide stock solution was serially diluted with nitrogen-bubbled sea water to produce sulfide dialysate solutions of 0.05, 0.10, 0.25, 0.50, and 1.00mM H_2S . Haemolymph samples were dialyzed for 24 hours at 4°C, extracted from dialysis cassettes, and fixed as above.

Fixed haemolymph samples were stored at room temperature for several weeks, then analyzed for sulfide content using a protocol modified from cline (1969). Briefly, 100 μ l of fixed sample was combined with 900 μ l distilled, deionized water, 20 μ l DPP solution (0.48 g N,N-dimethyl-p-phenylene diamine dihydrochloride in 100ml 6M HCl) and 20 μ l ferric chloride solution (1.6 g $FeCl_3$ in 100ml 6M HCl). After 60 min., samples were spun for 5 min. at low speed to remove large particles, and the absorbance of the supernatant was measured at 670 nm on a UV-vis diode-array spectrophotometer (HP 8452A). Sulfide concentrations in the fixed haemolymph samples were calculated from a standard curve, based on solutions of known sulfide concentration

processed simultaneously with haemolymph samples.

3. Result

3.1 Haemolymph Sulfide Binding

Sulfide concentrations in haemolymph from freshly-collected *Calyptogena soyoe* ranged from 0.1–0.4mM, with a mean of 0.21mM ($n=7$). Sulfide in the lamellibrachiid worm was comparable, near 0.25mM. *Bathymodiolus platifrons* had less sulfide in the haemolymph (0.001, 0.016, and 0.049mM in three individuals tested).

Sulfide binding experiments demonstrate that *Calyptogena soyoe* and *Lamellibrachia* sp. contain haemolymph which binds sulfide at high affinity, whereas haemolymph of *Bathymodiolus platifrons* exhibited no such affinity (Fig. 1). Haemolymph of *C. soyoe* showed the greatest affinity for sulfide among these species, with 1.5 ± 0.5 (s.d., $n=7$) mM bound at 0.1 to 0.25 mM H_2S , or a 6 to 8-fold concentration factor. At lower ambient sulfide levels (0.05mM H_2S), haemolymph concentrations also decreased, but were 20-fold greater than environmental levels. The affinity of lamellibrachiid haemo-




Fig. 1 Plot of haemolymph binding affinity from *Calyptogena soyoe*, *Bathymodiolus platifrons*, and an undescribed species of *Lamellibrachia*, from the Hatsushima cold seep site in Sagami Bay, Japan. Data represent sulfide concentrations in dialysed blood samples incubated for 24h at specified ambient sulfide levels.

lymph was slightly lower, with 0.83 mM bound at 0.25 mM H₂S (n=1), and a concentration factor less than 10 for all dialysate levels. In both species, H₂S in haemolymph appeared to approach an asymptotic limit with increasing dialysate H₂S concentration suggesting saturation of the binding moiety at external concentrations well above 1 mM H₂S. Linear regressions performed on double-reciprocal plots of these data suggest that haemolymph approaches saturation in *C. soyoae* near 2 mM H₂S, and 1.5 mM H₂S for *Lamellibrachia* sp. Sulfide levels in *Bathymodiolus platifrons* haemolymph were lower than dialysate concentrations in all trials.

4. Discussion

4.1 Role of Haemolymph Sulfide Binding Affinity

The ability of chemosynthetic species to bind sulfide very likely influences both their distribution among sites and their metabolic potential. Because delivery of sulfide from the seep pore waters to intracellular bacterial symbionts is facilitated by haemolymph, the capacity of haemolymph to bind sulfide is key in relating environmental chemistry to organism function (e.g., metabolism and growth). In vesicomyid clams and vestimentiferan worms, sulfide binding ability plays three potential roles. First, it allows the host to concentrate sulfide, thereby providing bacterial symbionts with intracellular sulfide levels that are considerably higher than exist externally. Second, it protects the host from the toxic effects of sulfide by sequestering sulfide from the host tissue in a "bound" form. Third, it provides a sulfide-rich reservoir that can be utilized by the symbionts during periods of low environmental sulfide availability.

Patterns of sulfide binding affinity and species-habitat associations observed for vesicomyid bivalves in Monterey Bay indicate that binding affinity is highest for species inhabiting low sulfide habitats, and vice versa. *Calyptogena pacifica* is most abundant at seeps with low interstitial sulfide levels (<0.1 mM H₂S; Barry *et al.* in press b), and can concentrate sulfide in haemolymph well over 10X ambient levels, and perhaps over 100X ambient concentrations. In contrast, *C. kilmeri*, which thrives in seeps with sulfide levels up to 100 times higher (>5 mM H₂S) than sites inhabited by *C. pacifica*, has a much

lower binding affinity for sulfide (Kochavar *et al.*, ms).

Calyptogena soyoae inhabits seeps in Sagami Bay with sulfide concentrations near 0.5 mM H₂S in the top 20 cm of sediment (Hashimoto *et al.* 1995). This level is intermediate between the low sulfide levels (<0.1 mM H₂S) typical of seeps in Monterey Bay dominated by *C. pacifica*, and the high sulfide seeps (>5 mM H₂S) in which *C. kilmeri* dominates. Based on these ambient pore fluid sulfide levels, we expected the haemolymph binding affinity of *C. soyoae* to be intermediate between the two Monterey vesicomyids. Surprisingly, the sulfide binding affinity of haemolymph from *C. soyoae* is unexpectedly high, and even exceeds *C. pacifica* at intermediate levels of dialysate.

Assuming that binding affinity for sulfide at least partially determines the levels of sulfide available to bacterial symbionts for vesicomyids, several potential explanations can be proposed for the divergent pattern of sulfide binding and habitat association among vesicomyids from Monterey and Sagami Bay cold seeps. First, symbionts are known to differ among species (Distal *et al.*, 1994), and may have vastly differing requirements for sulfide, thereby decoupling patterns of sulfide binding and habitat sulfide levels among species. Presently, we have no data to evaluate this hypothesis. Second, although sulfide binding affinity may partially determine habitat tolerance among species (e.g. *C. pacifica* and *C. soyoae* tolerate low sulfide environments), differences in binding affinity may be related to the combined effects of habitat tolerance and metabolic demand. Thus, high binding affinity may enable species increase the rate of sulfide transport to symbionts, as well as maintaining high haemolymph sulfide levels, thereby supporting higher metabolic activity (e.g. higher growth; see below). Finally, high binding affinity may result from a combination of factors, including the ability to tolerate spatial or temporal changes in sulfide concentration, or both. Thus, environmental sulfide levels near *C. soyoae* may be more temporally variable than in Monterey Bay, requiring a high binding affinity to support symbionts during periods of low environmental sulfide availability, but also a high capacity for tolerance during periods of high sulfide levels.

This last hypothesis is supported by our finding of low

sulfide in haemolymph from freshly-collected *C. soyoae*, suggesting that environmental levels may be low. Analyses of haemolymph from freshly-collected vesicomid clams and lamellibrachiid worms suggest that sulfide levels to which these animals are exposed are low-on the order of 0.1 mM H₂S. These findings are consistent with measurements of very low sulfide concentrations in haemolymph of freshly-collected *Bathymodiolus platifrons*. Congeneric mytilids found in hydrothermal vents and hydrocarbon seep settings, harboring either thiotrophic (*Bathymodiolus thermophilus*) or methanotrophic (Seep Mytilid Ia) symbionts, have not been found to concentrate sulfide from the external environment (Childress and Fisher, 1992; Kochevar *et al.*, in prep); thus their haemolymph sulfide concentrations approximate *in situ* sulfide levels.

4.2 Physiological Ecology of Thiotrophic Metazoan Symbioses

The combined effects of haemolymph sulfide binding affinity, symbiont sulfide requirements, and interstitial sulfide levels appear to play a large role in the ecological distribution of chemosynthetic species. High haemolymph sulfide affinity allows *Calyptogena pacifica* to inhabit seeps with very low environmental sulfide levels, presumably by enabling this species to elevate intracellular sulfide levels well above external levels at the point of delivery (i.e., the symbionts). Furthermore, the apparent de-coupling of carbon uptake from sulfide uptake (Kochevar *et al.*, in prep.) suggests that the haemolymph also serves an important reservoir function, buffering symbionts from fluctuations in external sulfide levels. However, the high affinity of *C. pacifica* haemolymph also results in saturation of the binding moiety at relatively low external sulfide levels, making this species vulnerable to sulfide toxicity in high sulfide habitats. *C. kilmeri*, on the other hand, exhibits low haemolymph sulfide affinity. This may reflect a relaxation of the need to maintain a reservoir of sulfide for the symbionts, suggesting a steady supply of sulfide in the environment; and may also provide a higher level of sulfide tolerance than that of *C. pacifica*, thus allowing it to thrive in habitats with much higher sulfide levels. Thus, there appear to be significant physiological trade-offs between sulfide tolerance and energetic uptake (i.e. bac-

terial productivity).

Because vesicomid clams are found intimately associated with sulfide-rich substrate, they have often been characterized as "bridging the gap" between reducing and oxidizing conditions in a spatial sense. This is in contrast to vestimentiferan worms, which are adapted for exposure to oxidizing and reducing conditions which alternate temporally on very rapid time scales. However, as our knowledge about vesicomids increases, particularly in the context of cold seep habitats, it is clear that physiological adaptations, as well as species distribution, are likely the product of both spatial and temporal variability in sulfide availability. Furthermore, the co-occurrence of vesicomid clams, vestimentiferan worms, and mytilid mussels at Sagami Bay cold seeps provides a unique model system in which to refine our understanding of the functioning of these three groups of organisms.

Studies of Monterey Bay and Sagami Bay cold seeps to date have provided much useful information about water chemistry, animal physiology, and community structure. The striking similarities among the various cold seep settings suggest that ongoing comparative studies will help lead to a broader understanding of these habitats worldwide. To better understand Monterey Bay and Sagami Bay cold seep communities, and to facilitate comparisons among them, we must begin to integrate detailed single-time-point observations with longer-term measurements which will provide a temporal context in which to understand patterns of change in water chemistry and animal distribution. Furthermore, ongoing developments at MBARI and JAMSTEC in *in situ* monitoring technology hold great promise for furthering our understanding the physical and chemical processes which underlie observed patterns in physiology and ecology of chemosynthetic fauna.

Acknowledgements

We are indebted to the pilots and support crews of the *R/V* Pt. Lobos and Ventana ROV and the *R/V* "Natsushima" and "Shinkai 2000" submersible. Support for the project was provided by JAMSTEC, Monterey Bay Aquarium Research Institute, and the United States-Japan Submersible Science Program, funded by NOAA.

References

- Aharon, P. (1994) : Geology and biology of modern and ancient submarine hydrocarbon seeps and vents : An Introduction. *Geo-Marine Letters*, 14 (2/3), 69–73.
- Barry, J. P., Greene, H. G., Orange, D. L., Baxter, C. H., Robison, B. H., Kochevar, R. E., Nybakken, J. W., Reed, D. L., and C. M. McHugh (1996a) : Biologic and Geologic Characteristics of Cold Seeps in Monterey Bay, California. *Deep-Sea Research* (in press).
- Barry, J. P., R. E. Kochevar, and C. H. Baxter (1996b) : The influence of pore-water chemistry and physiology in the distribution of vesicomyid clams at cold seeps in Monterey Bay : Implications for patterns of chemosynthetic community organization. *Limnology and Oceanography*, (in press).
- Beauchamp, B., H. R. Krouse, J. C. Harrison, W. W. Nassichuk, and L. S. Eluik (1989) : Cretaceous Cold-Seep Communities and Methane-Derived Carbonates in the Canadian Arctic. *Science*, 244, 53–56.
- Boss, K. J. and R. D. Turner (1980) : The giant white clam from the Galapagos Rift, *Calyptogena magnifica* species novum. *Malacologia*, 20, 161–194.
- Boulégué, J., E. L. Benedetti, D. Drou, A. Moriotti and R. Létolle (1987) : Geochemical and biogeochemical observations on the biological communities associated with fluid venting in Nankai trough and Japan trench subduction zones. *Earth and Planetary Science Letters*, 83, 343–355.
- Cadet, J.-P., K. Kobayashi, J. Aubouin, Boulégué, C. Deplus, J. Dubois, R. von Huene, L. Jolivet, T. Kanazawa, J. Kashara, K. Koizumi, S. Lallemand, Y. Nakamura, G. Pautot, K. Suyehiro, S. Tani, H. Tokuyama, and Y. Yamazaki (1987) : The Japan Trench and its juncture with the Kurile Trench : cruise results of the Kaiko project, Leg 3. *Earth and Planetary Science Letters*, 83, 268–284.
- Campbell, K. A. and D. J. Bottjer (1993) : Fossil Cold Seeps, *National Geographic Research Expeditions*, 9 (3), 326–343.
- Childress, J. J. and C. R. Fisher (1992) : The biology of hydrothermal vent animals : physiology, biochemistry, and autotrophic symbioses. *Oceanogr. Mar. Biol. Annu. Rev.*, 30, 337–441.
- Clari, P., L. Fornara, B. Ricci, and G. M. Zuppi (1994) : Methane-derived carbonates and chemoautotrophic communities of Piedmont (Miocene, northern Italy) : An update. *Geo-Marine Letters*, 14(2/3), 201–209.
- Cline, J. D. (1969) : Spectrophotometric determination of hydrogen sulfide in natural waters. *Limnology and Oceanography*, 14, 454–458.
- Dando, P. R., M. C. Austen, R. A. Burke Jr., M. A. Kendall, M. C. Kennicutt II, A. G. Judd, D. C. Moore, S. C. M. O'Hara, R. Schmaljohann, A. J. Southward (1991) : Ecology of a North Sea pockmark with an active methane seep. *Marine Ecology Progress Series*, 70, 49–63.
- Distel, D. L., H. Felbeck, and C. M. Cavanaugh (1994) : Evidence for phylogenetic congruence among sulfur-oxidizing chemoautotrophic bacterial endosymbionts and their bivalve hosts. *J. Molecular Evolution*, 38, 533–542.
- Embley, R. W., S. L. Eittreim, C. H. McHugh, W. R. Normark, G. H. Rau, B. Hecker, A. E. DeBevoise, H. G. Greene, W. B. F. Ryan, C. Harrold, and C. Baxter (1990) : Geological setting of chemosynthetic communities in the Monterey Fan Valley System. *Deep-Sea Research*, 37 (11), 1651–1677.
- Greene, H. G., J. P. Barry, J. Hashimoto, Y. Fujiwara, R. E. Kochevar, and B. H. Robison (1997) : A submersible-based Comparison of Cold-seep Regions in Sagami and Monterey Bays. *JAMSTEC J. Deep Sea Res.*, 12, 403–423.
- Hashimoto, J., Ohta, S., Fujikura, K., Fujiwara, Y., and S. Sukizaki (1995) : Life habit of vesicomyid clam, *Calyptogena soyoe*, and hydrogen sulfide concentration in interstitial waters in Sagami Bay, Japan. *J. Ocean.*, 51, 341–350.
- Hashimoto, J. T., Ohta, T., Tanaka, S., Matsuzawa, and H. Hotta (1987) : Surveys of the deep sea communities dominated by the giant clam, *Calyptogena soyoe* along the slope foot of Hatsushima Island, Sagami Bay. *JAMSTEC J. Deep Sea Res.*, 3, 37–50.
- Hashimoto, J., S. Ohta, T. Tanaka, H. Hotta, S. Matsuzawa, and H. Sakai (1989) : Deepsea communities dominated by the giant clam, *Calyptogena soy-*

- oe, along the slope foot of Hatsushima Island, Sagami Bay, Central Japan. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 71, 179–192.
- Hecker, B. (1985) : Fauna from a cold sulfur-seep in the Gulf of Mexico : comparison with hydrothermal vent communities and evolutionary implications. *Bulletin, Biology Society of Washington*, 6, 465–474.
- Jollivet, D., J.-C. Faugeres, R. Griboulard, D. D. Desbruyeres, and G. Blanc (1990) : Composition and spatial organization of a cold seep community on the South Barbados accretionary prism : tectonic, geochemical and sedimentary context. *Progress in Oceanography*, 24, 25–45.
- Laubier, L., S. Ohta and M. Sibuit (1986) : Découverte de communautés animales profondes durant la campagne franco-japonaise Kaiko de plongées dans les fosses de subduction autour du Japon. *C. R. Academy of Sciences of Paris*, t. 303, serie II, 2, 25–29.
- Le Pichon, X., T. Liyama, H. Chamley, J. Charvet, M. Faure, H. Fujimoto, T. Furuta, Y. Ida, H. Kagimi, S. Lallemant, J. Leggett, A. Murata, H. Okada, C. Rangin, V. Renard, A. Taira, and H. Tokuyama (1987) : The eastern and western ends of Nankai trough : results of box 5 and box 7 Kaiko survey. *Earth and Planetary Science Letters*, 83, 199–213.
- Kennicutt II, M. C., J. M. Brooks, R. R. Bidigare, and G. J. Denoux (1988) : Gulf of Mexico hydrocarbon seep communities-1. Regional distribution of hydrocarbon seepage and associated fauna. *Deep-Sea Research*, 35A, 1639–1651.
- Kennicutt II, M. C., J. M. Brooks, R. R. Bidigare, R. R. Fay, T. L. Wade, and T. J. McDonald (1985) : Vent-type taxa in a hydrocarbon seep region on the Louisiana Slope. *Nature*, 317 (6035), 35–353.
- Kennicutt II, M. C., J. M. Brooks, S. A. Macko, R. R. Bidigare, S. J. McDonald, and D. Adkison (1989) : An upper slope “cold” seep community : northern California. *Limnology and Oceanography*, 34, 635–640.
- Kochevar, R. E. and J. P. Barry (in preparation) : Physiology of vesicomyid clams from Monterey Bay cold seeps.
- Lonsdale, P. (1979) : A deep hydrothermal site on a strike-slip fault. *Nature*, 281, 531–535.
- Masuzawa, T., N. Handa, H. Kitagawa and M. Kusakabe (1992) : Sulfate reduction using methane in sediments beneath a bathyal “cold seep” giant clam community off Hatsushima Island, Sagami Bay, Japan. *Earth and Planetary Science Letters*, 110, 39–50.
- Masuzawa, T., T. Nakatsuka, and N. Handa (1995) : Geochemistry of pore water from a bathyal *Calyptogena* community off Hatsushima Island, Sagami Bay, Japan. Pages 407 to 421 in : *Biogeochemical Processes and Ocean Flux in the Western Pacific*, Eds, H. Sakai and Y. Nozaki, Terra Sci. Publ Company, Tokyo.
- Naka, J., K. Fujikura, and J. Hashimoto (1991) : New geological interpretation of *Calyptogena* colony off Hatsushima Island, Sagami Bay. *JAMSTEC J. Deep Sea Res.*, 7, 1–5.
- Ohta, S. and L. Laubier (1987) : Deep biological communities in the subduction zone of Japan from bottom photographs during ‘Nautil’ dives in the Kaiko project. *Earth and Planetary Science Letters*, 83, 329–342.
- Paull, C. K., Hecker, B., Commeau, C., Freeman-Lynde, R. P. Neumann, C., Corso, W. P., Golubic, G., Hook, J., Sikes, E., and J. Curaray (1984) : Biological communities at Florida Escarpment resemble hydrothermal vent communities. *Science*, 226, 965–967.
- Suess, E., Carson, S. D. Ritger, J. C. Moore, M. L. Jones, L. D. Kulm and G. R. Cochrane (1985) : Biological communities at vent sites along the subduction zone off Oregon. *Bulletin Biology Society of Washington*, 6, 475–484.
- von Bitter, P. H., S. D. Scott, and P. E. Schenk (1990) : Early carboniferous low-temperature hydrothermal vent communities from Newfoundland. *Nature*, 344 (6262), 145–148.
- von Bitter, P. H., S. D. Scott, and P. E. Schenk (1992) : Chemosynthesis : an alternate hypothesis for carboniferous biotas in bryozoan/microbial mounds, Newfoundland, Canada. *Palaios*, 7 (4), 466–484.

(Manuscript received 7, July 1997)