

## 4D-VAR Ocean State Estimation

December 1, 2015

### ■ Name of data set and the abbreviated name

**Name:** Estimated state of ocean for climate research

**Abbreviation:** ESTOC

### ■ Data Overview

Our 4D-VAR data synthesis system is developed as a part of the Japan Agency for Marine-Earth Science and Technology (JAMSTEC)-Kyoto University collaborative program (known as "the K7 consortium"). The Ocean General Circulation Model is version 3 of the GFDL Modular Ocean Model (MOM3), which is equipped with several sophisticated schemes; e.g., Noh scheme for mixed layer physics, the Gent and McWilliams (GM) scheme for isopycnal mixing, and quicker advection scheme. The horizontal resolution is  $1^\circ$  in both latitude and longitude, with 46 vertical levels spaced from 10 m near the sea surface to 400 m at the bottom. This model is better able to reproduce ocean circulation processes and is expected to form a platform suitable for the use of the 4D-VAR adjoint model.

4D-VAR adjoint data assimilation approach is applied. The adjoint codes of the OGCM were obtained using the Tangent linear and Adjoint Model Compiler (TAMC) and the Transformation of Algorithms in Fortran (TAF). In the 4D-VAR approach, optimized 4-dimensional datasets are sought by minimizing a cost function.

Physical parameters:

The assimilated elements in this study are temperature and salinity from ENSEMBLES (EN3) dataset which was quality controlled using a comprehensive set of objective checks developed at the Hadley Centre of the UK Meteorological Office. This dataset is largely composed of observations from the World Ocean Database 2005 and supplemented by data from the GTSP (Global Temperature and Salinity Profile Program) and Argo autonomous profiling floats. In addition of EN3 dataset, recent independent MIRAI RV profiles are simultaneously synthesized. Sea-surface dynamic-height anomaly data derived from high-precision multi-satellite altimetry products distributed by Aviso is also incorporated. All observational data were averaged onto  $1^\circ$  by  $1^\circ$  bins and then compiled as series of 10-day means for the surface data and monthly means for the subsurface data. The control variables are surface fluxes (for net-heat, fresh water, and momentum) and oceanic initial conditions. The assimilation window is 55 years during 1957-2011.

Biogeochemical parameters:

The synthesis of available observations and a pelagic ecosystem model based on nitrogen cycle produces a dynamically self-consistent dataset. Optimized 4-dimensional datasets are sought by minimizing a cost function on the basis of Green's function approach. The assimilated elements are the climatological monthly mean nitrate from WOA05, monthly mean ocean color data from SeaWiFS, and annual mean chlorophyll-a from WOA98 as detritus.

## ■ Dataset release date

December 1, 2015

## ■ Corresponding contributor

Nozomi Sugiura

Address: Research and Development Center for Global Change,

Japan Agency for Marine-Earth Science and Technology

2-15 Natsushima-cho, Yokosuka, Kanagawa

237-0061, Japan

Telephone: +81-46-867-9054, FAX: +81-46-867-9835

E-mail: smasuda at jamstec.go.jp

## ■ Terms and Conditions

The copyright of data of this site belongs to JAMSTEC.

The terms and conditions are as follows:

1. JAMSTEC shall not be responsible for any loss, damages and troubles you make either directly or indirectly by using downloaded data of this site.
2. You are not permitted to manipulate, redistribute, deliver, or sell downloaded data of this site to public.
3. Uses of downloaded data for scientific or educational purposes  
Subject to the following A) to D), using downloaded data for nonprofit scientific or educational purposes is free of charge.
  - A) If you use downloaded data, you are requested to express that JAMSTEC data is used in your paper or document, etc. Please follow the citation format in the description for the dataset (i.e. this PDF file) included in downloaded data.
  - B) If you publish a paper in a journal or make a presentation at an academic conference using downloaded data, please let us know the information such as offprint or URL of abstract, etc.
  - C) If you share downloaded data within your laboratory or project, please let us know the information such as name of laboratory or project.
  - D) If you hand out downloaded data at university lectures or seminars, etc., please let us know the information such as name of university and course.
4. Uses of downloaded data for non-scientific or non-educational purposes  
If you use downloaded data for non-scientific or non-educational purposes, such as commercial purpose, etc., please inform us in advance and follow our instructions.

Contact: estoc-admin at jamstec.go.jp

## ■ References to be cited

Please cite as follows:

Osafune, S., S. Masuda, N. Sugiura, T. Doi (2015) Evaluation of the applicability of the Estimated Ocean State for Climate Research (ESTOC) dataset, *Geophys. Res. Lett.*, 42, 12, 4903–4911.

Doi, T., S. Osafune, N. Sugiura, S. Kouketsu, A. Murata, S. Masuda, and T. Toyoda (2015) Multi-decadal change in the dissolved inorganic carbon in a long-term ocean state estimation, *Journal of Advances in Modeling Earth Systems*, in press.

## ■ Available Products

Variables, abbreviations	Potential temperature [°C], Salinity [PSU], Horizontal velocity u[m/s] v[m/s], Surface heat flux [cal/m <sup>2</sup> /s], Surface freshwater flux [m/s], Wind stress $\tau_x$ [N/m <sup>2</sup> ] $\tau_y$ [N/m <sup>2</sup> ], Nitrate [ $\mu$ mol/L], Phytoplankton [ $\mu$ mol/L], Detritus [ $\mu$ mol/L], Zooplankton [ $\mu$ mol/L], Dissolved inorganic carbon [ $\mu$ mol/kg], Dissolved oxygen [ $\mu$ mol/L],	tmp sal vel shf sff tau no3 pht det zoo dic oxy
Region	Quasi-global (75°S-80°N)	
Resolution	Horizontal 1°x1°, Vertical 46 levels	
Period	1957-2011 (Ver. 02c)	
File format	Monthly data in netcdf format: “k7oda_[XXX]_[YYYY][MM]00_[VVV].nc” where “[XXX]” is model variable, “[YYYY]” year, “[MM]” month, and “[VVV]” version, respectively.	

## ■ Reference URL

- K7 Database  
<http://www.jamstec.go.jp/frcgc/k7-dbase2/eng/index.html>
- K7 Ocean State Estimate based on 4D-VAR Ocean Data Assimilation  
[http://www.jamstec.go.jp/frcgc/k7-dbase2/eng/datadoc/k7ra\\_ocean.html](http://www.jamstec.go.jp/frcgc/k7-dbase2/eng/datadoc/k7ra_ocean.html)
- Data and Sample Research System for Whole Cruise Information in JAMSTEC (DARWIN)  
<http://www.godac.jamstec.go.jp/darwin/e>
- Japan Argo Delayed-mode Data base  
[http://www.jamstec.go.jp/ARGO/argo\\_web/argo/?lang=en](http://www.jamstec.go.jp/ARGO/argo_web/argo/?lang=en)